

# Median preserving aggregation functions

Miguel Couceiro

LORIA

(CNRS, INRIA Nancy G.-E., Université de Lorraine)

Joint work with Jean-Luc Marichal and Bruno Teheux

# Outline

## **I. Brief overview on aggregation theory :**

- I.1. Aggregation functions : motivation
- I.2. An impossibility result : Arrow's theorem

## **II. Aggregation over median algebras :**

- II.1. Median algebras : motivation and examples
- II.2. Conservative median algebras
- II.3. Median preserving aggregation : An Arrow-like theorem

## I.1 Aggregation functions

**Traditionally** : an **aggregation function** is a mapping  $F: X^n \rightarrow X$  **s.t.**

- $X$  is a linear order with bottom 0 and top 1
- $F$  preserves 0 and 1 and the order of  $X$

**Typical examples** : Weighted means, Choquet and Sugeno integrals ...

**Main Idea** : Aggregation procedure  $x_1, \dots, x_n \in X \rightarrow F(x_1, \dots, x_n) \in Y$

**Application** : Preference modelling (MCDA) ...

**Main Problems** :

- Classify and axiomatise aggregation procedures
- *Explicitly describe procedures with desired properties*
- Computational aspects

## I.1 Aggregation functions

**Traditionally** : an **aggregation function** is a mapping  $F: X^n \rightarrow X$  **s.t.**

- $X$  is a linear order with bottom 0 and top 1
- $F$  preserves 0 and 1 and the order of  $X$

**Typical examples** : Weighted means, Choquet and Sugeno integrals ...

**Main Idea** : Aggregation procedure  $x_1, \dots, x_n \in X \rightarrow F(x_1, \dots, x_n) \in Y$

**Application** : Preference modelling (MCDA) ...

**Main Problems** :

- Classify and axiomatise aggregation procedures
- *Explicitly describe procedures with desired properties*
- Computational aspects

## I.1 Aggregation functions

**Traditionally** : an **aggregation function** is a mapping  $F: X^n \rightarrow X$  **s.t.**

- $X$  is a linear order with bottom 0 and top 1
- $F$  preserves 0 and 1 and the order of  $X$

**Typical examples** : Weighted means, Choquet and Sugeno integrals ...

**Main Idea** : Aggregation procedure  $x_1, \dots, x_n \in X \rightarrow F(x_1, \dots, x_n) \in Y$

**Application** : Preference modelling (MCDA) ...

**Main Problems** :

- Classify and axiomatise aggregation procedures
- *Explicitly describe procedures with desired properties*
- Computational aspects

## I.2. An impossibility result : Arrow's theorem

**Setting** : Aggregation of rankings (social well-fare function)

- $n$  voters, a set  $A$  of outcomes and the set of linear orderings  $L(A)$
- $F: L(A)^n \rightarrow L(A)$  procedure that merges rankings  $R_1, \dots, R_n$  into a single one

$$R_1, \dots, R_n \implies R_T = F(R_1, \dots, R_n)$$

- Some reasonable properties in this setting :
  1. **Unanimity or Pareto efficiency** : if  $aR_i b$  for all  $i \in [n]$ , then  $aR_T b$
  2. **Independence of irrelevant alternatives** : if  $a$  and  $b$  have the same order in  $R_i$  and  $S_j$  for all  $i \in [n]$ , then  $a$  and  $b$  have the same order in  $R_T$  and  $S_T$
  3. **Non-dictatorship** : There is no  $i \in [n]$  s.t. for all  $R_1, \dots, R_n \in L(A)$

$$F(R_1, \dots, R_n) = R_i$$

**Arrow's Theorem** : There is no well-fare function satisfying these conditions !

## I.2. An impossibility result : Arrow's theorem

**Setting :** Aggregation of rankings (social well-fare function)

- $n$  voters, a set  $A$  of outcomes and the set of linear orderings  $L(A)$
- $F: L(A)^n \rightarrow L(A)$  procedure that merges rankings  $R_1, \dots, R_n$  into a single one

$$R_1, \dots, R_n \implies R_T = F(R_1, \dots, R_n)$$

- Some reasonable properties in this setting :
  1. **Unanimity or Pareto efficiency :** if  $aR_i b$  for all  $i \in [n]$ , **then**  $aR_T b$
  2. **Independence of irrelevant alternatives :** if  $a$  and  $b$  have the same order in  $R_i$  and  $S_i$  for all  $i \in [n]$ , **then**  $a$  and  $b$  have the same order in  $R_T$  and  $S_T$
  3. **Non-dictatorship :** There is no  $i \in [n]$  **s.t.** for all  $R_1, \dots, R_n \in L(A)$

$$F(R_1, \dots, R_n) = R_i$$

**Arrow's Theorem :** There is no well-fare function satisfying these conditions !

## I.2. An impossibility result : Arrow's theorem

**Setting :** Aggregation of rankings (social well-fare function)

- $n$  voters, a set  $A$  of outcomes and the set of linear orderings  $L(A)$
- $F: L(A)^n \rightarrow L(A)$  procedure that merges rankings  $R_1, \dots, R_n$  into a single one

$$R_1, \dots, R_n \implies R_T = F(R_1, \dots, R_n)$$

- Some reasonable properties in this setting :
  1. **Unanimity or Pareto efficiency :** if  $aR_i b$  for all  $i \in [n]$ , **then**  $aR_T b$
  2. **Independence of irrelevant alternatives :** if  $a$  and  $b$  have the same order in  $R_i$  and  $S_i$  for all  $i \in [n]$ , **then**  $a$  and  $b$  have the same order in  $R_T$  and  $S_T$
  3. **Non-dictatorship :** There is no  $i \in [n]$  **s.t.** for all  $R_1, \dots, R_n \in L(A)$

$$F(R_1, \dots, R_n) = R_i$$

**Arrow's Theorem :** There is no well-fare function satisfying these conditions !



## II.1. Median algebras : motivation

**Median operations** appear in several structures pertaining to decision making :

- **Linear orders** : “in betweenness”
- **Distributive lattices** :  $\mathbf{m}(x, y, z) = (x \wedge y) \vee (y \wedge z) \vee (z \wedge x)$

**Theorem** : A function  $f: X^n \rightarrow X$  is a lattice polynomial function (Sugeno integral) iff

$$f(\mathbf{x}) = \mathbf{m}(f(\mathbf{x}_k^0), x_k, f(\mathbf{x}_k^1)) \quad \text{for every } \mathbf{x} \in X^n, k \in [n]$$

## II.1. Median algebras : motivation

**Median operations** appear in several structures pertaining to decision making :

- **Linear orders** : “in betweenness”
- **Distributive lattices** :  $\mathbf{m}(x, y, z) = (x \wedge y) \vee (y \wedge z) \vee (z \wedge x)$

**Theorem** : A function  $f: X^n \rightarrow X$  is a lattice polynomial function (Sugeno integral) **iff**

$$f(\mathbf{x}) = \mathbf{m}(f(\mathbf{x}_k^0), x_k, f(\mathbf{x}_k^1)) \quad \text{for every } \mathbf{x} \in X^n, k \in [n]$$

## II.1. Median algebras : definition and examples

**Median algebra** : Structure  $\mathbf{A} = (A, \mathbf{m})$  where  $\mathbf{m} : A^3 \rightarrow A$  (median) verifies

$$\mathbf{m}(x, x, y) = x$$

$$\mathbf{m}(x, y, z) = \mathbf{m}(y, x, z) = \mathbf{m}(y, z, x)$$

$$\mathbf{m}(\mathbf{m}(x, y, z), t, u) = \mathbf{m}(x, \mathbf{m}(y, t, u), \mathbf{m}(z, t, u))$$

Other known median algebras :

- **Median semilattices** : For  $a \in A$ , set  $x \leq_a y \iff \mathbf{m}(a, x, y) = x$
- **Median graphs** : For all  $x, y, z$ , there is a unique  $w$  in the shortest paths

**NB1** : Every median semilattice (with finite intervals) has a median Hasse diag.

**NB2** : Every median graph is the Hasse diagram of a median semilattice

**References** : Barthélemy-Leclerc-Monjardet'86, Bandelt'83, Isbell'80, Avann'61, ...

**Generalisations** : Bandelt-Meletiou'92, Barthélemy-Janowitz'91, Bandelt'90, ...

## II.1. Median algebras : definition and examples

**Median algebra :** Structure  $\mathbf{A} = (A, \mathbf{m})$  where  $\mathbf{m}: A^3 \rightarrow A$  (median) verifies

$$\begin{aligned} \mathbf{m}(x, x, y) &= x \\ \mathbf{m}(x, y, z) &= \mathbf{m}(y, x, z) = \mathbf{m}(y, z, x) \\ \mathbf{m}(\mathbf{m}(x, y, z), t, u) &= \mathbf{m}(x, \mathbf{m}(y, t, u), \mathbf{m}(z, t, u)) \end{aligned}$$

**Other known median algebras :**

- **Median semilattices :** For  $a \in A$ , set  $x \leq_a y \iff \mathbf{m}(a, x, y) = x$
- **Median graphs :** For all  $x, y, z$ , there is a unique  $w$  in the shortest paths

**NB1 :** Every median semilattice (with finite intervals) has a median Hasse diag.

**NB2 :** Every median graph is the Hasse diagram of a median semilattice

**References :** Barthélemy-Leclerc-Monjardet'86, Bandelt'83, Isbell'80, Avann'61, ...

**Generalisations :** Bandelt-Meletiou'92, Barthélemy-Janowitz'91, Bandelt'90, ...

## II.2. Conservative median algebras

**Conservative median algebra :** If  $m(x, y, z) \in \{x, y, z\}$ ,  $x, y, z \in A$

**Social choice motivation :** the median candidate is one of the candidates

**Problem :** How do they look like ?

## II.2. Conservative median algebras

**Conservative median algebra :** If  $m(x, y, z) \in \{x, y, z\}$ ,  $x, y, z \in A$

**Social choice motivation :** the median candidate is one of the candidates

**Problem :** How do they look like ?

## II.2. Conservative median algebras

**Conservative median algebra :** If  $m(x, y, z) \in \{x, y, z\}$ ,  $x, y, z \in A$


**Social choice motivation :** the median candidate is one of the candidates

**Problem :** How do they look like ?

## Representation of conservative median algebras

**Theorem :** Let  $\mathbf{A}$  be a median algebra with  $|A| \geq 5$ . T.F.A.E.

- (i)  $\mathbf{A}$  is conservative.
- (ii) There is an  $a \in A$  and lower bounded chains  $\mathbf{C}_0$  and  $\mathbf{C}_1$  such that  $\langle A, \leq_a \rangle$  is isomorphic to  $\mathbf{C}_0 \perp \mathbf{C}_1$ .
- (iii) For every  $a \in A$ , there are lower bounded chains  $\mathbf{C}_0$  and  $\mathbf{C}_1$  such that  $\langle A, \leq_a \rangle$  is isomorphic to  $\mathbf{C}_0 \perp \mathbf{C}_1$ .
- (iv) For every  $a \in A$  the ordered set  $\langle A, \leq_a \rangle$  does not contain a copy of the poset



**Open problem :** Representation of arbitrary median algebras




## II.3. Median preserving aggregation

**Idea :** Score of a median profile is the median of the scores of the profiles


**Problem :** Aggregation functions  $f: X^n \rightarrow Y$  that preserve medians :

$$f(\mathbf{m}(\mathbf{x}, \mathbf{y}, \mathbf{z})) = \mathbf{m}(f(\mathbf{x}), f(\mathbf{y}), f(\mathbf{z})),$$

**Remark :** median preserving maps are not necessarily order-preserving (reversing) !



An order-preserving map that is not median preserving



A median preserving map that is not order-preserving (or reversing)


## II.3. Median preserving aggregation

**Idea :** Score of a median profile is the median of the scores of the profiles


**Problem :** Aggregation functions  $f: X^n \rightarrow Y$  that preserve medians :

$$f(\mathbf{m}(\mathbf{x}, \mathbf{y}, \mathbf{z})) = \mathbf{m}(f(\mathbf{x}), f(\mathbf{y}), f(\mathbf{z})),$$

**Remark :** median preserving maps are not necessarily order-preserving (reversing) !



An order-preserving map that is not median preserving



A median preserving map that is not order-preserving (or reversing)

## Characterization of median preserving maps

**NB :** Every conservative median algebra  $\mathbf{A}$  can be thought of as a chain  $\mathbf{C}(\mathbf{A})$

**Theorem :** Let  $\mathbf{A}, \mathbf{B}$  be conservative median algebras with  $\geq 5$  elements. T.F.A.E. :

- (i)  $f : \mathbf{A} \rightarrow \mathbf{B}$  is a median preserving map
- (ii) the induced map  $f' : \mathbf{C}(\mathbf{A}) \rightarrow \mathbf{C}(\mathbf{B})$  is order-preserving or order-reversing

**Problem :** How to lift it to  $f : \mathbf{A}^n \rightarrow \mathbf{B}$

## Characterization of median preserving maps

**NB :** Every conservative median algebra  $\mathbf{A}$  can be thought of as a chain  $\mathbf{C}(\mathbf{A})$

**Theorem :** Let  $\mathbf{A}, \mathbf{B}$  be conservative median algebras with  $\geq 5$  elements. T.F.A.E. :

- (i)  $f : \mathbf{A} \rightarrow \mathbf{B}$  is a median preserving map
- (ii) the induced map  $f' : \mathbf{C}(\mathbf{A}) \rightarrow \mathbf{C}(\mathbf{B})$  is order-preserving or order-reversing

**Problem :** How to lift it to  $f : \mathbf{A}^n \rightarrow \mathbf{B}$

## Characterization of median preserving maps

**NB :** Every conservative median algebra  $\mathbf{A}$  can be thought of as a chain  $\mathbf{C}(\mathbf{A})$

**Theorem :** Let  $\mathbf{A}, \mathbf{B}$  be conservative median algebras with  $\geq 5$  elements. T.F.A.E. :

- (i)  $f : \mathbf{A} \rightarrow \mathbf{B}$  is a median preserving map
- (ii) the induced map  $f' : \mathbf{C}(\mathbf{A}) \rightarrow \mathbf{C}(\mathbf{B})$  is order-preserving or order-reversing

**Problem :** How to lift it to  $f : \mathbf{A}^n \rightarrow \mathbf{B}$

## Back to aggregation functions...

**Theorem :** Let  $\mathbf{A} = \mathbf{C}_1 \times \cdots \times \mathbf{C}_n$  and  $\mathbf{B} = \mathbf{D}_1 \times \cdots \times \mathbf{D}_k$  be products of chains. T.F.A.E. :

- (i)  $f : \mathbf{A} \rightarrow \mathbf{B}$  is median preserving
- (ii) there exist  $\sigma : [k] \rightarrow [n]$  and order-preserving or order-reversing maps

$$f_i : \mathbf{C}_{\sigma(i)} \rightarrow \mathbf{D}_i \quad \text{for } i \in [k] \quad \text{s.t. } f(\mathbf{x}) = (f_1(x_{\sigma(1)}), \dots, f_k(x_{\sigma(k)}))$$

**Corollary :** Let  $\mathbf{C}_1, \dots, \mathbf{C}_n$  and  $\mathbf{D}$  (in part.,  $k = 1$ ) be chains. T.F.A.E. :

- (i)  $f : \mathbf{C}_1 \times \cdots \times \mathbf{C}_n \rightarrow \mathbf{D}$  is median preserving
- (ii) there is  $j \in [n]$  and order-preserving or reversing map  $g : \mathbf{C}_j \rightarrow \mathbf{D}$  s.t.  $f = g \circ \pi_j$

**Consequence :** Arrow-like theorem over median algebras

*Aggregation procedures that preserve medians are dictatorial !*

## Back to aggregation functions...

**Theorem :** Let  $\mathbf{A} = \mathbf{C}_1 \times \cdots \times \mathbf{C}_n$  and  $\mathbf{B} = \mathbf{D}_1 \times \cdots \times \mathbf{D}_k$  be products of chains. T.F.A.E. :

- (i)  $f : \mathbf{A} \rightarrow \mathbf{B}$  is median preserving
- (ii) there exist  $\sigma : [k] \rightarrow [n]$  and order-preserving or order-reversing maps

$$f_i : \mathbf{C}_{\sigma(i)} \rightarrow \mathbf{D}_i \quad \text{for } i \in [k] \quad \text{s.t. } f(\mathbf{x}) = (f_1(x_{\sigma(1)}), \dots, f_k(x_{\sigma(k)}))$$

**Corollary :** Let  $\mathbf{C}_1, \dots, \mathbf{C}_n$  and  $\mathbf{D}$  (in part.,  $k = 1$ ) be chains. T.F.A.E. :

- (i)  $f : \mathbf{C}_1 \times \cdots \times \mathbf{C}_n \rightarrow \mathbf{D}$  is median preserving
- (ii) there is  $j \in [n]$  and order-preserving or reversing map  $g : \mathbf{C}_j \rightarrow \mathbf{D}$  s.t.  $f = g \circ \pi_j$

**Consequence :** Arrow-like theorem over median algebras

*Aggregation procedures that preserve medians are dictatorial !*

## Back to aggregation functions...

**Theorem :** Let  $\mathbf{A} = \mathbf{C}_1 \times \cdots \times \mathbf{C}_n$  and  $\mathbf{B} = \mathbf{D}_1 \times \cdots \times \mathbf{D}_k$  be products of chains. T.F.A.E. :

- (i)  $f : \mathbf{A} \rightarrow \mathbf{B}$  is median preserving
- (ii) there exist  $\sigma : [k] \rightarrow [n]$  and order-preserving or order-reversing maps

$$f_i : \mathbf{C}_{\sigma(i)} \rightarrow \mathbf{D}_i \quad \text{for } i \in [k] \quad \text{s.t. } f(\mathbf{x}) = (f_1(x_{\sigma(1)}), \dots, f_k(x_{\sigma(k)}))$$

**Corollary :** Let  $\mathbf{C}_1, \dots, \mathbf{C}_n$  and  $\mathbf{D}$  (in part.,  $k = 1$ ) be chains. T.F.A.E. :

- (i)  $f : \mathbf{C}_1 \times \cdots \times \mathbf{C}_n \rightarrow \mathbf{D}$  is median preserving
- (ii) there is  $j \in [n]$  and order-preserving or reversing map  $g : \mathbf{C}_j \rightarrow \mathbf{D}$  s.t.  $f = g \circ \pi_j$

**Consequence :** Arrow-like theorem over median algebras

*Aggregation procedures that preserve medians are dictatorial !*



*Merci de votre attention !*

*Thank you for your attention !*

*Obrigado pela vossa atenção !*

*Grazie mille per la vostra attenzione !*